



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Substituting  $\tan \alpha$  for  $m$ , we get

$$\frac{\pi}{2} \log \frac{1 + \sin \alpha}{1 - \sin \alpha} = \frac{\pi}{2} \log \frac{(1 + \sin \alpha)^2}{1 - \sin^2 \alpha} = \frac{\pi}{2} \log \left( \frac{1 + \sin \alpha}{\cos \alpha} \right)^2 = \pi \log (\tan \alpha + \sec \alpha),$$

which result is quite as great as that given in the proposed problem.

---

### MECHANICS.

---

188. Proposed by H. L. ORCHARD, M. A., B. Sc.

Spherical bubbles are rising in water. Find the relation between radius and velocity.

No solution has been received.

189 (Incorrectly numbered 190). Proposed by DR. L. E. DICKSON, The University of Chicago, Chicago, Ill.

Give the axiomatic principle of Physics which is equivalent to the theorem on the compound of two circles ("Graphical Methods in Trigonometry," MONTHLY, June-July, 1905).

No solution has been received.

190 (Incorrectly numbered 191). Proposed by J. EDWARD SANDERS, Reinersville, Ohio.

A pole hinged at the bottom leans against the mid-point of a smooth rope suspended from two supports of equal height. Determine the position of equilibrium.

Solution by G. W. GREENWOOD, M. A., Dunbar, Pa.

Assuming that the figure is symmetrical with respect to the plane bisecting perpendicularly the segment determined by the two points of support, the tensions of the portions of the rope resolved parallel to the pole are equal in magnitude and direction, and must therefore be zero since the rope is smooth. Hence, the pole rests in a position perpendicular to the plane determined by the portions of the rope.

191 (Incorrectly numbered 192). Proposed by REV. J. H. MEYER, S. J., College of Sacred Heart, Augusta, Ga.

Find the velocity of a planet at a given point in its orbit.

Solution by G. W. GREENWOOD, M. A., Dunbar, Pa.

In the case of any central orbit we have  $v = h/p \dots (1)$ , where  $p$  is the perpendicular from the pole upon the tangent at the point, and  $h$  is a constant. Also,

$$\frac{1}{p^2} = u^2 + \left( \frac{du}{d\theta} \right)^2$$

and the equation of an ellipse, referred to a focus, is  $lu = 1 + e \cos \theta$ , where  $l$  is the semi-latus rectum.